

# GTC OSIRIS SPECTROSCOPIC IDENTIFICATION OF A FAINT L SUBDWARF IN THE UKIRT INFRARED DEEP SKY SURVEY

N. LODIEU<sup>1,2</sup>, M. R. ZAPATERO OSORIO<sup>3</sup>, E. L. MARTÍN<sup>3</sup>, E. SOLANO<sup>3</sup>, M. ABERASTURI<sup>3</sup>

*Draft version December 17, 2009*

## ABSTRACT

We present the discovery of an L subdwarf in 234 deg<sup>2</sup> common to the UK Infrared Telescope (UKIRT) Infrared Deep Sky Survey Large Area Survey Data Release 2 and the Sloan Digital Sky Survey Data Release 3. This is the fifth L subdwarf announced to date, the first one identified in the UKIRT Infrared Deep Sky Survey, and the faintest known. The blue optical and near-infrared colors of ULAS J135058.86+081506.8 and its overall spectra energy distribution are similar to the known mid-L subdwarfs. Low-resolution optical (700–1000 nm) spectroscopy with the Optical System for Imaging and low Resolution Integrated Spectroscopy spectrograph on the 10.4 m Gran Telescopio de Canarias reveals that ULAS J135058.86+081506.8 exhibits a strong KI pressure-broadened line at 770 nm and a red slope longward of 800 nm, features characteristics of L-type dwarfs. From direct comparison with the four known L subdwarfs, we estimate its spectral type to be sdL4–sdL6 and derive a distance in the interval 94–170 pc. We provide a rough estimate of the space density for mid-L subdwarfs of  $1.5 \times 10^{-4}$  pc<sup>-3</sup>.

*Subject headings:* infrared: stars — stars: Population II — subdwarfs — techniques: photometric — techniques: spectroscopic

## 1. INTRODUCTION

Objects cooler than M dwarfs have been classified into two spectral classes: L and T (Martín et al. 1999; Kirkpatrick et al. 1999; Geballe et al. 2002; Burgasser et al. 2006). Hundreds of them are now known as a result of searches of L and T dwarfs in large scale surveys<sup>4</sup>. However, at lower metallicity, only four L subdwarfs have been announced to date: 2MASS J053253.46+824646.5 (2MASS0532; sdL7; Burgasser et al. 2003), 2MASS J162620.34+392519.0 (2MASS1626; sdL4; Burgasser 2004), 2MASS 061640.06–640719.4 (2MASS0616; sdL5; Cushing et al. 2009), and SDSS J125637.13–022452.4 (SDSS1256; sdL3.5; Sivarani et al. 2009; Burgasser et al. 2009). The first three were discovered serendipitously in the Two Micron All Sky Survey (2MASS; Cutri et al. 2003; Skrutskie et al. 2006) archive while the last was found during a search for ultracool subdwarfs in the Second Data Release (DR) of the Sloan Digital Sky Survey (SDSS; York et al. 2000; Abazajian et al. 2005) spectroscopic database. Increasing the number of ultracool subdwarfs is essential to study the chemistry in cool atmospheres with low metal content, the role of metallicity in the shape of the initial mass function (Salpeter 1955; Miller & Scalo 1979; Scalo 1986), and the impact of chemical composition on the properties of binary stars (Riaz et al. 2008; Jao et al. 2008; Lodieu et al. 2009b).

The UKIRT Deep Infrared Sky Survey (UKIDSS; Lawrence et al. 2007) uses the Wide Field Camera (WF-

CAM; Casali et al. 2007) installed on the UK InfraRed Telescope (UKIRT) and the Mauna Kea Observatory (Tokunaga et al. 2002) photometric system described in Hewett et al. (2006). The pipeline processing is described in Irwin et al. (in preparation)<sup>5</sup> and the WFCAM Science Archive in Hambly et al. (2008). The Large Area Survey (hereafter LAS), one of the UKIDSS components, will image  $\sim 3800$  deg<sup>2</sup> in four filters (*YJHK*) down to  $J = 19.6$  mag with a significant overlap with SDSS, hence providing spectral energy distributions from 0.3 to 2.5  $\mu$ m for millions of sources.

In this letter, we present the discovery of the first L subdwarf identified in 234 deg<sup>2</sup> common to the UKIDSS LAS DR2 (Warren et al. 2007) and the SDSS DR3 (Abazajian et al. 2005). In Section 2, we describe the photometric and proper motion selections to identify ultracool subdwarfs in UKIDSS and SDSS. In Section 3, we present the spectroscopic follow-up observations conducted with the Optical System for Imaging and low Resolution Integrated Spectroscopy (OSIRIS) spectrograph mounted on the new 10.4 m Gran Telescopio de Canarias (GTC) telescope in La Palma (Canary Islands). In Section 4, we discuss the photometric and spectral characteristics of the new object, infer its distance, and compute its tangential velocity. Finally, we conclude with a discussion on the expected number of subdwarfs that could be identified at the completion of UKIDSS.

## 2. SAMPLE SELECTION

We have initiated a photometric and proper motion search for ultracool (spectral types later than M7) subdwarfs by cross-correlating the UKIDSS and SDSS databases. In the LAS, we have requested only point sources (`mergedClass` = -1) and good quality detec-

Electronic address: nlodieu@iac.es

<sup>1</sup> Instituto de Astrofísica de Canarias, C/ Vía Láctea s/n, E-38205 La Laguna, Tenerife, Spain

<sup>2</sup> Departamento de Astrofísica, Universidad de La Laguna, E-38205 La Laguna, Tenerife, Spain

<sup>3</sup> Centro de Astrobiología (CSIC/INTA), 28850 Torrejón de Ardoz, Madrid, Spain

<sup>4</sup> See DwarfArchives.org for the full list of L and T dwarfs

<sup>5</sup> Extensive details on the data reduction of the UKIDSS images is available at <http://casu.ast.cam.ac.uk/surveys-projects/wfcam/technical>

tions ( $\text{ppErrBits} \leq 256$ ). We have employed the following photometric criteria:  $Y - J = 0.3\text{--}0.75$  mag,  $J - K \leq 0.7$  mag because of the strong collision-induced  $\text{H}_2$  absorption beyond  $2\mu\text{m}$  (Linsky 1969), and  $z - J \leq 2.5$  mag because the optical-to-infrared colors of subdwarfs are expected to be bluer than their solar metallicity counterparts (Section 4; Figure 1). We have also imposed constraints on the LAS detections, requesting  $J = 14\text{--}18.5$  mag and photometric errors less than 0.2 mag in  $Y$  and  $J$  and better than 0.25 mag in  $H$  and  $K$ . The bright limit in  $J$  was set to place our candidates well above the saturation of the UKIDSS LAS. In addition, we have requested SDSS detections below the  $5\sigma$  limits in  $u$  ( $\geq 22.3$  mag) and  $g$  ( $\geq 23.3$  mag). Besides the photometric criteria, we have imposed a lower limit of  $0.18 \text{ arcsec yr}^{-1}$  on the proper motion (Luyten 1980), which is measured as the difference between UKIDSS and SDSS coordinates.

This query returned seven photometric candidates. The photometry of six candidates suggest that they are late-M subdwarfs. Spectroscopic follow-up of these six sources is currently underway. However, one object, ULAS J135058.86+081506.8 (hereafter ULAS1350), is the only source with a negative  $J - H$  color and among the faintest ones displaying optical and near-infrared colors similar to those of 2MASS1626 (sdL4; Burgasser 2004), the second L subdwarf known at the time of our discovery. Table 1 provides the catalogued photometry of ULAS1350 and two other known L subdwarfs as well as a field L5 dwarf for comparison. Therefore, we prioritized this candidate for optical spectroscopy that we finally secured with the GTC in 2009. From the SDSS/UKIDSS cross-match, we inferred a total proper motion of  $0.280 \pm 0.025 \text{ arcsec yr}^{-1}$  thanks to the  $\sim 3$  yr baseline between both observations (April 2003 and July 2006). The typical uncertainty on the difference between the UKIDSS and SDSS astrometry is  $0.025 \text{ arcsec yr}^{-1}$  for sources brighter than  $J = 18.5$  mag (see analysis in Dye et al. (2006) and Lodieu et al. (2009a)). This proper motion is lower than the motion of the other four L subdwarfs currently known. ULAS1350 is also fainter, suggesting that it lies at a larger distance. Its reduced proper motion ( $H_i = 23.45$  mag) is similar to 2MASS1626 ( $H_i = 23.34$  mag) and SDSS1256 ( $H_i = 23.41$  mag), supporting its possible low-metallicity status.

### 3. SPECTROSCOPIC OBSERVATIONS

The 10.4 m GTC started operations in 2009 March at the Observatory del Roque de Los Muchachos (La Palma, Canary Islands). It is currently equipped with one of its Day One instruments, the OSIRIS instrument (Cepa et al. 2000). The OSIRIS spectrograph consists of two  $2048 \times 4096$  Marconi CCD42-82 with an 8 arcsec gap between them and operates at optical wavelengths, from 365 to 1000 nm. The unvignetted instrument field of view is  $7 \times 7$  arcmin with a pixel scale is  $0.125 \text{ arcsec}$ . This Letter reports on one of the first scientific results obtained with GTC/OSIRIS after the gamma-ray burst circular of Castro-Tirado et al. (2009) and the observations of SGR 0418+5729 by Mignani et al. (2009).

We have carried out low-resolution ( $R \sim 515$  at  $8500\text{\AA}$ ) spectroscopy of ULAS1350 with the R500R grism available on GTC/OSIRIS and a slit width of 1 arcsec, projecting onto a full-width-half-maximum of seven pixels

onto the detector, yielding a 480–1000 nm wavelength coverage and a nominal dispersion of  $0.244 \text{ nm pixel}^{-1}$ . The observations were obtained in service mode by the GTC staff on 2009 April 30. Spectra were taken at parallactic angle. Weather conditions were spectroscopic and seeing around  $1.0 \text{ arcsec}$ . The total exposure time was 2100 s in one single exposure. Skyflats and bias frames were observed on 2009 April 4.

All data were reduced under the IRAF<sup>6</sup> environment, which includes bias subtraction, and flat-field correction using sky-flat images. The individual spectrum was optimally extracted and the Earth sky emission lines were removed with the APALL task by choosing the appropriate aperture width and background intervals. Wavelength calibration (in the air system) was performed with an accuracy of 0.05 nm using the internal arc lamp lines of Argon and Neon, which were acquired at the end of the night in which ULAS1350 was observed. To correct for the instrumental response we used twilight sky-flat images (taken with the same instrumental configuration as our target) that recorded the solar spectrum. The closest observation of a spectrophotometric standard with the R500R grating was carried out on 2009 April 4 (HZ 44; Oke 1990), the same night as the skyflats. We have used it to confirm that the correction from instrumental response using the twilight flat images is reliable in the wavelength range 700–900 nm, where this star has spectrophotometric information in the IRAF database. Fringing is negligible ( $\leq 1\%$ ) shortwards of 900 nm. The extracted spectrum of ULAS1350 has a signal-to-noise ratio of around 3 at around 920–925 nm. The usable optical (700–1000 nm) spectrum of ULAS1350 smoothed with a boxcar of 51 pixels and a signal-to-noise ratio of  $\sim 20$  is displayed in Figure 2 and compared to the spectra (resolution degraded to that of ULAS1350 and similarly smoothed) of the four known L subdwarfs from the literature.

## 4. ANALYSIS

### 4.1. Photometry

From the optical and near-infrared photometry alone, ULAS1350 is unlikely to be a solar-metallicity object because it shows discrepant colors compared to late-M and L dwarfs (Table 1). Its  $J - K$  color of  $-0.02 \pm 0.16$  mag is typical of T dwarfs and much bluer than solar-metallicity field L dwarfs discovered by 2MASS and SDSS ( $J - K \geq 1.2$  mag; Hawley et al. 2002). Moreover, the  $z - J$  and  $i - z$  are also much bluer with values of  $1.55 \pm 0.07$  and  $1.74 \pm 0.11$  mag compared to  $\sim 2.5\text{--}3.3$  mag and  $> 2.0$  mag, for mid-L dwarfs with solar composition, respectively.

Figure 3 shows the optical to near-infrared spectral energy distribution (SED) of ULAS1350 compared to 2MASS1626 (sdL4; Burgasser 2004) and a solar-metallicity L5 dwarf, SDSS J144600.60+002452.0 (Geballe et al. 2002). The SED of ULAS1350 clearly suggests a sub-solar metallicity especially in the near-infrared, as also inferred from Figure 1 and Table 1. To convert observed magnitudes into physical fluxes we have used the zero point fluxes corresponding to each SDSS

<sup>6</sup> IRAF is distributed by National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation.

and UKIDSS passband defined in Hewett et al. (2006). We have flux calibrated the GTC data by integrating the observed spectrum convolved with the response curves of the *i* and *z* filters (Fukugita et al. 1996).

#### 4.2. Spectral features

The GTC spectrum confirms the cool and low-luminosity atmosphere of our candidate, for which we determined  $\text{sdL5} \pm 1$  after comparison to the four known bright L subdwarfs (Fig. 2). Indeed, the spectrum of ULAS1350 clearly exhibits a strong pressure-broadened KI band at  $\sim 770$  nm (Figure 2) as well as a red slope longwards of 800 nm, features typical of early to mid-L field dwarfs. We also detect the hydride bands of CrH ( $\sim 860$  nm) and FeH ( $\sim 870$  nm and  $\sim 990$  nm), typically stronger features in the spectra of lower metallicity stars (Mould 1976). From Figure 2, our spectrum shows a decreasing flux at wavelengths below 730 nm, a feature that is also shared by the known  $\text{sdL3.5}$ – $\text{sdL5}$  objects. It is likely due to absorption by TiO gas, which is usually stronger in the subdwarfs relative to solar L dwarfs (probably an effect due to inhibited dust formation in low-metallicity atmospheres (e.g. Burgasser et al. 2003; Reiners et al. 2006). While Burgasser et al. (2007) proposed a recipe for assigning spectral types to L subdwarfs by comparing with known solar-metallicity L dwarfs, no set of spectral standards is currently defined due to the few L subdwarfs known, so we consider our spectral type as tentative.

#### 4.3. Distance and tangential velocity

We have considered the absolute magnitude versus spectral type relations given by Cushing et al. (2009) to estimate the distance of ULAS1350 as no L5 subdwarf with known trigonometric parallax exist. Assuming a spectral type of  $\text{sdL5}$  and an uncertainty of one subtype, we derive  $M_J = 12.715$  mag ( $12.402$ – $13.028$  mag), yielding a mean distance of 111 pc with a possible range from 96 to 128 pc. Using the relations for *H* and *K*, we find a distance 25–30% larger than for the *J*-band relation. The uncertainty on the distance due to the relations of Cushing et al. (2009) is 13–14 pc, implying that the uncertainty on the distance is dominated by the error on the spectral classification. To account for both uncertainties, we added the errors in quadrature, leading to a spectroscopic distance of  $140^{+30}_{-46}$  pc for ULAS1350.

Combining the proper motion ( $0.28$  arcsec  $\text{yr}^{-1}$ ) and the distance (140 pc) derived above, we derive a tangential velocity of  $186^{+40}_{-61}$   $\text{km.s}^{-1}$  where the error bars account for the distance uncertainty. On the one hand, these values are  $\sim 2$ – $4$  times larger than the mean values of tangential velocity reported for L dwarfs in the solar neighborhood (Vrba et al. 2004; Faherty et al. 2009). On the other hand, the transverse velocity of ULAS1350 is quite similar to the tangential velocities of the ultracool subdwarfs shown in Table 2 by Schilbach et al. (2009). This indicates that ULAS1350 likely exhibits halo kinematics. As inferred from Figure 1, state-of-the-art models predict a metallicity between  $[\text{M}/\text{H}] = -0.5$  and  $-1.0$  dex for ULAS1350 although current models do not reproduce accurately the color trends of field dwarfs mainly because condensate clouds are not included, resulting in offsets between predicted and observed colors (Figure 1;

Burgasser et al. (2009)).

### 5. DISCUSSION

ULAS1350 was discovered in a 234 square area common to the SDSS and UKIDSS surveys. Our search was limited to *J* magnitudes brighter than 18.5. Therefore, we can provide an estimate of the density of L5 subdwarfs in the surveyed area. At the distance of 140 pc, the explored volume is approximately  $6600 \text{ pc}^{-3}$ , implying a rough space density of  $1.5 \times 10^{-4}$   $\text{sdL5s}$  per cubic parsecs. The density could, however, be larger as we have six other photometric candidates identified in the same area and we have not yet explored the faint end of the LAS. This discovery could also be fortuitous, yielding an overestimate of the space density. Moreover, ULAS1350 could be a multiple system leading to the sampling of a larger volume. The inferred density is  $\sim 20$  times lower than the space density of solar-type field L dwarfs (lower limit of  $3.8 \times 10^{-3} \text{ pc}^{-3}$ ; Cruz et al. 2007). It is also 10 times higher than the density of M subdwarfs for  $M_V = 5$ – $10$  ( $1$ – $3 \times 10^{-5} \text{ pc}^{-3}$ ; Digby et al. 2003); we note however that our estimation is obtained for fainter and cooler objects, which may have smaller masses. This might indicate that the subdwarf mass function (in a linear scale) is flat or slowly rising toward the Hydrogen burning-mass limit. We remark that this is a tentative suggestion since it is based on the finding of one L5 subdwarf for which distance is not known precisely. If we extrapolate our findings to the final area imaged by the LAS with SDSS photometry ( $\sim 3000 \text{ deg}^2$ ), we would expect roughly 13 mid-L subdwarfs. This tentative number shows that our search may open new prospects of designing an accurate spectral classification in the L dwarf regime including metallicity as a variable parameter (Kirkpatrick 2005). With this aim, we have conducted a cross-match of UKIDSS DR6 and SDSS DR7 following a Virtual Observatory methodology whose results will be presented in a forthcoming paper.

N.L. was funded by the Ramón y Cajal fellowship number 08-303-01-02. This work was partially funded by Spanish Ministry of Science grant number AYA2007-67458 and the Consolider-Ingenio 2010 Program grant CSD2006-00070: “First Science with the GTC (<http://www.iac.es/consolider-ingenio-gtc>)”. NL thanks numerous discussions on subdwarfs with Richard Jameson during his post-doc in Leicester. We thank the GTC staff for their help as well as Adam Burgasser and Mike Cushing for providing the optical spectra of the known L subdwarfs used in this Letter. We thank the anonymous referee for his/her prompt and useful report.

Based on observations made with the Gran Telescopio Canarias (GTC), installed in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias, in the island of La Palma.

The United Kingdom Infrared Telescope is operated by the Joint Astronomy Centre on behalf of the U.K. Science Technology and Facility Council.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western

TABLE 1  
PHOTOMETRY AND SPECTRAL TYPE OF ULAS J135058.86+081506.8  
ALONG WITH TWO COMPARISON L SUBDWARFS AND A FIELD L DWARF.

	ULAS1350 <sup>a</sup>	SDSS1256 <sup>b</sup>	2MASS1626 <sup>b</sup>	SDSS1446 <sup>a,c</sup>
R.A.	13:50:58.86	12:56:37.10	16:26:20.34	14:46:00.60
Decl.	+08:15:06.8	-02:24:52.5	+39:25:19.0	+00:24:52.0
$i - z$	1.74±0.11	1.70±0.02	1.74±0.01	2.27±0.06
$z - J$	1.55±0.07	1.61±0.02	1.71±0.01	2.92±0.03
$J - K$	-0.02±0.16	-0.10±0.02	-0.03±0.01	1.66±0.01
$r$	24.30±0.66	21.80±0.11	20.61±0.03	23.19±0.24
$i$	21.22±0.09	19.39±0.02	17.89±0.01	20.77±0.05
$z$	19.48±0.06	17.68±0.02	16.15±0.01	18.50±0.03
$Y$	18.60±0.05	—	—	16.90±0.01
$J$	17.94±0.04	16.16±0.01	14.43±0.01	15.58±0.01
$H$	18.08±0.10	16.06±0.01	14.46±0.01	14.66±0.01
$K$	17.98±0.15	16.06±0.02	14.46±0.01	13.92±0.01
SpT	sdL5±1.0	sdL3.5±0.5	sdL4±0.5	dL5±0.5

<sup>a</sup> Optical photometry from SDSS DR7 (AB mag) and near-infrared photometry from UKIDSS LAS DR6 (Vega system)

<sup>b</sup> Optical photometry from SDSS DR7 (AB mag) and near-infrared photometry from Schilbach et al. (2009)

<sup>c</sup> Original discovery from Geballe et al. (2002)

Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the

Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

## REFERENCES

- Abazajian K., et al. 2005, *AJ*, 129, 1755  
 Baraffe I., Chabrier G., Allard F., Hauschildt P. H., 1997, *A&A*, 327, 1054  
 Burgasser A. J., 2004, *ApJL*, 614, L73  
 Burgasser A. J., Cruz K. L., Kirkpatrick J. D., 2007, *ApJ*, 657, 494  
 Burgasser A. J., Geballe T. R., Leggett S. K., Kirkpatrick J. D., Golimowski D. A., 2006, *ApJ*, 637, 1067  
 Burgasser A. J., et al. 2003, *ApJ*, 592, 1186  
 Burgasser A. J., Witte S., Helling C., Sanderson R. E., Bochanski J. J., Hauschildt P. H., 2009, *ApJ*, 697, 148  
 Casali M., et al. 2007, *A&A*, 467, 777  
 Castro-Tirado A. J., et al. 2009, GRB Coordinates Network, 9655, 1  
 Cepa J., et al. 2000, in M. Iye & A. F. Moorwood ed., Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series Vol. 4008 of Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, OSIRIS tunable imager and spectrograph. pp 623–631  
 Cruz K. L., et al. 2007, *AJ*, 133, 439  
 Cushing M. C., et al. 009, *ApJ*, 696, 986  
 Cutri R. M., et al. 2003, 2MASS All Sky Catalog of point sources, 2246  
 Digby A. P., Hambly N. C., Cooke J. A., Reid I. N., Cannon R. D., 2003, *MNRAS*, 344, 583  
 Dye S., et al. 2006, *MNRAS*, 372, 1227  
 Faherty J. K., Burgasser A. J., Cruz K. L., Shara M. M., Walter F. M., Gelino C. R., 2009, *AJ*, 137, 1  
 Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimasaku K., Schneider D. P., 1996, *AJ*, 111, 1748  
 Geballe T. R., et al. 2002, *ApJ*, 564, 466  
 Hambly N. C., et al. 2008, *MNRAS*, 384, 637  
 Hawley S. L., et al. 2002, *AJ*, 123, 3409  
 Hewett P. C., Warren S. J., Leggett S. K., Hodgkin S. T., 2006, *MNRAS*, 367, 454  
 Jao W.-C., Henry T. J., Beaulieu T. D., Subasavage J. P., 2008, *AJ*, 136, 840  
 Kirkpatrick J. D., 2005, *ARA&A*, 43, 195  
 Kirkpatrick J. D., et al. 1999, *ApJ*, 519, 802  
 Kirkpatrick J. D., et al. 2000, *AJ*, 120, 447  
 Lawrence A., et al. 2007, *MNRAS*, 379, 1599  
 Lépine S., Scholz R.-D., 2008, *ApJL*, 681, L33  
 Linsky J. L., 1969, *ApJ*, 156, 989  
 Lodieu N., Leggett S. K., Bergeron P., & Nitta A. 2009a, *ApJ*, 692, 1506  
 Lodieu N., Zapatero Osorio M. R., Martín E. L., 2009b, *A&A*, 499, 729  
 Luyten W. J., 1980, NLTT catalogue. Vol.3: 0 deg. to -10 deg.. Minneapolis: University of Minnesota, 1980  
 Martín E. L., Delfosse X., Basri G., Goldman B., Forveille T., Zapatero Osorio M. R., 1999, *AJ*, 118, 2466  
 Metchev S. A., Kirkpatrick J. D., Berriman G. B., Looper D., 2008, *ApJ*, 676, 1281  
 Mignani, R. P., et al. 2009, The Astronomer's Telegram, 2044, 1  
 Miller G. E., Scalo J. M., 1979, *ApJS*, 41, 513  
 Mould J. L., 1976, *ApJ*, 207, 535  
 Oke J. B., 1990, *AJ*, 99, 1621  
 Reiners A., Basri G., 2006, *AJ*, 131, 1806  
 Riaz B., Gizis J. E., Samaddar D., 2008, *ApJ*, 672, 1153  
 Salpeter E. E., 1955, *ApJ*, 121, 161  
 Scalo J. M., 1986, *Fundamentals of Cosmic Physics*, 11, 1  
 Schilbach E., Röser S., Scholz R., 2009, *A&A*, 493, L27  
 Scholz R.-D., Lodieu N., McCaughrean M. J., 2004, *A&A*, 428, L25  
 Sivarani T., Lépine S., Kembhavi A. K., Gupchup J., 2009, *ApJL*, 694, L140  
 Skrutskie M. F., et al. 2006, *AJ*, 131, 1163  
 Tokunaga A. T., Simons D. A., Vacca W. D., 2002, *PASP*, 114, 180  
 Vrba F. J., et al. 2004, *AJ*, 127, 2948  
 Warren S. J., et al. 2007, *arXiv:astro-ph/0703037*  
 West A. A., et al. 2008, *AJ*, 135, 785  
 York D. G., et al. 2000, *AJ*, 120, 1579




FIG. 1.—  $(i - J, J - K)$  diagram showing the location of two ultracool subdwarfs, SDSS J023557.61+010800.5 (esdM7) and SDSS J020533.75+123824.0 (esdM8.5) from Lépine & Scholz (2008) and three L subdwarfs as large dots: ULAS1350 (sdL5), 2MASS1626 (sdL4), and SDSS1256 (sdL3.5) with SDSS photometry. Photometric error bars have been added for ULAS1350 and SDSS J023557.61+010800.5; the others being smaller than the size of the symbols. Theoretical 10 Gyr tracks for different metallicities are also shown (Baraffe et al. 1997). The colors of M0–M9 dwarfs are overplotted as open triangles (West et al. 2008); L dwarfs lie outside this diagram because of their red  $J - K$  colors. A similar version of this plot was used by Scholz et al. (2004) and Burgasser et al. (2009).




FIG. 2.— Smoothed ( $R \sim 10$ ) optical spectrum of ULAS1350 from the GTC. The signal-to-noise ratio of the spectrum after smoothing is of the order of 20 over the 920–925 nm wavelength range. Overplotted are the optical spectra of the four known L subdwarfs: 2MASS0532 (Keck/LRIS; sdL7; Burgasser et al. 2003), 2MASS0616 (Gemini/GMOS; sdL5; Cushing et al. 2009), 2MASS1626 (Gemini/GMOS; sdL4; Burgasser et al. 2007), and SDSS1256 (Magellan/LDSS3; sdL3.5; Burgasser et al. 2009) degraded in wavelength resolution and smoothed in the same manner as ULAS1350. Spectra have been normalized at 8200Å and are shifted along the y-axis by 1.2 for clarity.




FIG. 3.— Observed SED of ULAS1350 formed by combining SDSS/UKIDSS photometry and the GTC spectrum (circles and solid line). For comparison purposes, the observed SEDs of the metal-depleted dwarf 2MASS1626 (sdL4; triangles and dashed line; optical and near-infrared photometry from SDSS and Schilbach et al. (2009), respectively) and the solar metallicity dwarf SDSS1446 (dL5; squares and dotted line; photometry from SDSS and UKIDSS) are also shown. While the slope in the visible wavelengths are rather similar for solar and metal-depleted mid-L dwarfs, the SED appears quite different in the near-infrared.